

**ARIZONA GAME AND FISH DEPARTMENT
HERITAGE DATA MANAGEMENT SYSTEM**

Plant Abstract

Element Code: PDAP0030M0

Data Sensitivity: Yes

CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE



ME: *Amsonia kearneyana*

COMMON NAME: Kearney's Blue Star, Kearney's blue-star, Kearney's bluestar, Kearney's Amsonia

SYNONYMS:

FAMILY: Apocynaceae

AUTHOR, PLACE OF PUBLICATION: R.E. Woodson, Ann. Missouri Bot. Gard. 15(4): 415. 1928.

TYPE LOCALITY: USA, Arizona, Pima County, south canyon (Baboquivari Mts.), 9 April 1928, F. Thackery #55, MBG.

TYPE SPECIMEN: Type: MO, F.A. Thackery, 9 April 1928, Arizona. Paratype: US 01367358, F.A. Thackery 2018, 24 Mar 1926, USA. Arizona. Baboquivari Mts., south canyon.

TAXONOMIC UNIQUENESS: The species *kearneyana* is 1 of 16 in the genus *Amsonia*, and 1 of 5 in the subgenus *Sphinctosiphon*, which also includes: *A. tharpii*, *A. jonesii*, *A. palmeri*, and *A. peeblesii*. *A. kearneyana* was considered as synonymous with *A. palmeri* (North American Flora 29:129), but Kearney et al. (1951) maintains *A. kearneyana* based on distinct characteristics of mature follicles. McLaughlin (1982) retains species based on geographic separation between species, larger corolla lobes, and stem pubescence. Not a synonym of *A. palmeri* as suggested by Woodson (1928). Should be retained as distinct species. McLaughlin (1982): A specimen of *A. palmeri* from the Rio Bavispe, northeastern Sonora, has some characteristics of *A. kearneyana*.

DESCRIPTION: Perennial, milky sap multi-stemmed herb up to 90 cm (35.4 in) tall, with a thickened woody root. Up to 50 pilose stems are usually clustered from the base, giving plant hemispherical shape; sparingly branched and densely pubescent. Alternate to subverticillate, oblong-lanceolate to lanceolate leaves (lower: 6-10 cm [2.4-4.0 in.] long and 11-17 mm broad, upper: 4-6 cm [1.6-2.4 in] long and 3-8 mm broad), with soft spreading hairs along the margins. The inflorescence is not too conspicuous. White flowers (with pale pinkish/bluish bottom) are borne in clusters at the ends of the inflorescence. The pubescent calyx is 3.0-5.0 mm long, while the salverform **corolla** (long tube **with short lobes** [2-4 mm long] on the end) is 12-15 mm (range 10-22 mm) long, broadest below the apex and constricted at the orifice. The fruit is a pair of follicles, 3.0-10.0 cm (1.2-4.0 in.) long, slightly constricted between seeds. Seeds are cylindrical, corky, 8-11 mm long and 3-4 mm broad. *A. kearneyana* has largest seed size for its sub-group within *Amsonia*.

AIDS TO IDENTIFICATION: *A. kearneyana* has pubescent foliage and calyx; large seed 3-4 mm wide and 8-11 mm long; short corolla lobes, 2-4 mm long. Not sympatric with any other *Amsonias*. Closest *Amsonia* is *A. palmeri* with very showy flowers, narrower leaves, and different pubescence. *A. grandiflora* typically occurs at slightly higher elevations in oak woodlands and is glabrous, its leaves are longer and narrower, its corolla is longer, and its fruits are terete and not constricted between seeds (Falk, Jenkins et al. 2001).

ILLUSTRATIONS: Line drawing. USFWS.

Color photo (Accessed 6/18/2003 from <http://arizonaes.fws.gov/images/Jrorabaugh/05-30-2000/PlantsB/Kearneys%20bluestar.jpg>)

Line drawing (*In* Falk, Jenkins et al. 2001)

Color photo of plant (F. Reichenbacher, *in* Falk, Jenkins et al. 2001).

Color photo of plant in habitat (J. Rorabaugh, *in* Falk, Jenkins et al. 2001).

Color photo (Kathy Rice CPC #119, Accessed 6/18/2003 from http://ridgwaydb.mobot.org/cpcweb/CPC_ProfileImage.asp?NF=119a)

Color photos of plant and habitat (Accessed 6/18/2003 from <http://genome-lab.ucdavis.edu/People/RickTopinka/kearneyana.htm>)

Color drawing (Accessed 6/18/2003 from <http://www.co.pima.az.us/cmo/sdcp/sdcp2/fsheets/kbs.html>)

TOTAL RANGE: Western slopes of the Baboquivari Mountains, Pima County, Arizona.

RANGE WITHIN ARIZONA: South and Sycamore canyons, Baboquivari Mountains, Pima County. Introduced into Brown Canyon, east side of the Baboquivari Mountains.

SPECIES BIOLOGY AND POPULATION TRENDS

GROWTH FORM: Herbaceous perennial.

PHENOLOGY: Flowers late April and May, fruiting June through August. Seeds sterile (Woodson 1928). However, McLaughlin (1982) states that sterility due to insect predation on embryo.

BIOLOGY: Insect predators attack embryo giving sterile appearing seeds. Responds to winter precipitation. Reichenbacher has found variability in flower shapes and sizes in plants from the introduction site, implying that the population has some genetic variability. Hawk moths may pollinate at night. As plants mature, become five to six feet across.

HABITAT: Dry, open, slopes (20-30 degrees) at 4,000-6,000 ft (1220-1830 m) elevation in Madrean evergreen woodlands/interior chaparral transition zone and on stable, partially-shaded, coarse alluvium along dry washes at 3,600-3,800 ft (1095-1160 m) elevation under deciduous riparian trees and shrubs in Sonoran desertscrub or desertscrub-grassland ecotone (Falk, Jenkins et al. 2001).

ELEVATION: 3,600 - 6,400 ft. (1097 - 1950 m).

EXPOSURE:

SUBSTRATE: Granitic alluvium.

PLANT COMMUNITY: “Mexican Blue Oak association, Sonoran Desertscrub, Semidesert Grassland plant communities, or a transition zone between the two” (Reichenbacher, 1993). Madrean evergreen woodland/interior chaparral transition zone to Sonoran desertscrub or desertscrub-grassland ecotone (Falk, Jenkins et al. 2001). Associated plants (Van Devender 1981b) include: *Prosopis velutina*, *Brumelia lanuginosa*, *Coursetia glandulosa*, *Baccharis sarothroides*, *Quercus oblongifolia*, *Dasyllirion wheeleri*, *Celtis reticulata*, *Juglans major*, *Forestiera*, *Cocculus diversifolius*, *Acacia greggii*, *Anisacanthus thurberi*, *Phacelia*, *Cryptantha*, *Crossosoma biglovii*, *Gossypium thurberi*, *Dodonaea viscosa*, *Vauquelinia pauciflora*, *Ptelea trifoliolata*, *Vitis arizonica*, and *Nicotiana trigonophylla*.

POPULATION TRENDS: Location of this plant forgotten between 1928 and mid-1970's. McLaughlin found plant in South Canyon (8 individuals in entire population). In 1987, Howell surveyed for additional locations but none were found. From 1987 to 1989, U.S. Fish and Wildlife Service contracted with Southwestern Field Biologists to transplant Arizona Sonoran Desert Museum seeds from South Canyon to establish a new population. In spring of 1988, about 90 plants were planted in Brown Canyon, which is now part of the Buenos Aires National Wildlife Refuge. In 1989, 110 plants were also planted in Brown Canyon. The introduced population in Brown Canyon (east side of Baboquivari Mountains) declined from approximately 130 to 35 plants (2/3 of the population) following a flood in 1990. (At the Bureau of Land Management, Safford District, Rare Plants Workshop, Frank Reichenbacher stated that the flood occurred in July, 1991. However, his report of 1991 gives the year as 1990).

The one native population consists of approximately 10-15 individuals (late 1980's). Native population in 1986 and 1987 was 8-12 individuals in field. Flooding occurred in 1988, 1989 and 1992. By November 1993, had 65 plants surviving. Low recruitment due partially to seed pods being collected for two years because insects were boring into pods and eating seeds. No young in South Canyon at all. The status of the Sycamore Canyon population is unknown.

SPECIES PROTECTION AND CONSERVATION

ENDANGERED SPECIES ACT STATUS: LE (USDI, FWS 1989)
[C1 USDI, FWS 1985]
[C1 USDI, FWS 1980]

STATE STATUS: Highly Safeguarded (ARS, ANPL 1999)
[Highly Safeguarded (ARS, ANPL 1993)]

OTHER STATUS: Not Forest Service Sensitive (USDA, FS
Region 3 1999)

[Forest Service Sensitive (USDA, FS
Region 3 1990)]

MANAGEMENT FACTORS: Extreme rarity, vulnerable canyon bottom habitat. Protection of populations from disturbance. Physical damage from livestock.

CONSERVATION MEASURES TAKEN: Artificial population established by transplants in Brown Canyon; Baboquivari protected area on Tohono O'odham Reservation.

SUGGESTED PROJECTS: Surveys should be done in the Coyote Mountains (S. Rutman 1994).

LAND MANAGEMENT/OWNERSHIP: BIA - Tohono O'odham Nation; BLM - Phoenix and Tucson Field Offices; Private. Introduction site on FWS - Buenos Aires National Wildlife Refuge.

SOURCES OF FURTHER INFORMATION

REFERENCES:

- Arizona Revised Statutes, 1993. Arizona Native Plant Law. Appendix A:1.
Arizona Revised Statutes, 1999. Arizona Native Plant Law. Appendix A:1.
Center for Plant Conservation (CPC) – National Collection of Endangered Plants. Plant Profile *Amsonia kearneyana*. Accessed 6/18/2003 from http://ridgwaydb.mobot.org/cpcweb/CPC_ViewProfile.asp?CPCNum=119.
Desert Botanical Garden (DBG). Web abstract *Amsonia kearneyana*, Kearney's Blue Star. http://67.29.153.101:28485/Collections/amsonia_kearneyana.html. Accessed: 6/18/2003.
Falk, M., P. Jenkins, et al; Arizona Rare Plant Committee. 2001. Arizona Rare Plant Guide. Published by a collaboration of agencies and organizations. U.S. Government Printing Office. Pages unnumbered.
Fletcher, R.A. 1979. *Amsonia kearneyana*. Provisional status report. Prepared for USDA-Forest Service, Region 3, Albuquerque, New Mexico.
Integrated Taxonomic Information System (ITIS). Retrieved 6/18 2003 from ITIS, <http://www.itis.usda.gov>.
Kearney, T.H., R.H. Peebles with collaborators. 1960. Arizona flora. Second edition with supplement by J.T. Howell, E. McClintock and collaborators. University of California Press, Berkeley. p.653.
Kelly, K. and J. McGinnis. 1994. Highly safeguarded protected native plants of Arizona. Arizona Department of Agriculture, Native Plant Protection Program, Phoenix, Arizona.
Lehr, J.H. 1978. A catalogue of the flora of Arizona. Desert Botanical Garden, Phoenix, Arizona. p. 119.
McLaughlin, S.P. 1982. A revision of the southwestern species of *Amsonia* (Apocynaceae). *Annals of Missouri Botanical Garden* 69:336-350.
Missouri Botanical Garden – TROPICOS, Nomenclatural Data Base. *Amsonia kearneyana* Woodson. http://mobot.mobot.org/cgi-bin/search_vast?onda=N01804525. Accessed: 6/18/2003.

- Phillips, B.G. and N. Brian. 1982. Status report on *Amsonia kearneyana* Woodson. For U.S. Fish and Wildlife Service, Albuquerque New Mexico. 10 pp.
- Reichenbacher, F.W, L.A. Clifford-Reichenbacher and J.D. Taiz. 1991. Transplantation and monitoring of Kearney's blue star, Brown Canyon, Baboquivari Mountains. Prepared for U.S. Fish and Wildlife Service, Ecological Services. Phoenix, Arizona. 35 pp.
- Reichenbacher, F.W. and J.L. Welch. 1993. Kearney's blue star (*Amsonia kearneyana*) recovery plan. Prepared for U.S. Fish and Wildlife Service, Ecological Services. Phoenix, Arizona. 25 pp.
- Reichenbacher, F.W. 1994. Bureau of Land Management, Safford District, Rare Plant Workshop. November 14-16. Tucson, Arizona.
- Shreve, F. and I.L. Wiggins. 1964. Vegetation and flora of the Sonoran Desert. Stanford University Press, Stanford, California. p.1099.
- USDA, Forest Service Region 3. 1990. Regional Forester's Sensitive Species List.
- USDA, Forest Service Region 3. 1999. Regional Forester's Sensitive Species List.
- USDA, NRCS. 2002. The PLANTS Database, Version 3.5 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.
- USDI, Fish and Wildlife Service. 1975. Threatened or Endangered Fauna or Flora: Review of Status of Vascular Plants and Determination of "Critical Habitat"; Notice of Review. Federal Register 40(127):27827.
- USDI, Fish and Wildlife Service. 1980. Endangered and Threatened Wildlife and Plants; Review of Plant Taxa for Listing as Endangered or Threatened Species. Notice of Review. Federal Register 45(242):82486.
- USDI, Fish and Wildlife Service. 1985. Endangered and Threatened Wildlife and Plants; Review of Plant Taxa for Listing as Endangered or Threatened Species. Notice of Review. Federal Register 50(188):00002.
- USDI, Fish and Wildlife Service. 1987. Endangered Species Technical Bulletin. Kearney's Blue-Star (*Amsonia kearneyana*). 12(8):5
- USDI, Fish and Wildlife Service. 1989. Endangered and Threatened Wildlife and Plants; Determination of *Amsonia kearneyana* to be an endangered species. Federal Register 54(12):2131-2134.
- U.S. National Herbarium Type Specimen Register (US). Department of Systematic Biology – Botany, Smithsonian Institution. Accessed 6/18/2003 from <http://rathbun.si.edu/botany/fullRecords.cfm?myFamily=>.
- Van Devender, T.R. 1981a. Status report: *Amsonia kearneyana* (Kearney's Blue Star) for the Arizona Natural Heritage Program. 2 pp.
- Van Devender, T.R. 1981b. Notes on *Amsonia kearneyana*. Separate pers. comm. with Steve McLaughlin and Barb Phillips (April 1981 survey notes).
- Woodson, R.E. Jr. 1928. Studies in the Apocynaceae. III, a monograph of the genus *Amsonia*. Annals of Missouri Botanical Garden. 14:379-434

MAJOR KNOWLEDGEABLE INDIVIDUALS:

Jefford Francisco - Tucson, Arizona.

Steve McLaughlin - University of Arizona, Office of Arid Lands Studies, Tucson, Arizona.

Barb Phillips - USFS Zone Botanist, Coconino, Kaibab and Prescott National Forests, Flagstaff, Arizona.

Frank Reichenbacher - Southwestern Field Biologists, Tucson, Arizona.

ADDITIONAL INFORMATION:

Revised: 1991-11-10 (PLW)
1991-12-04 (SR)
1994-12-07 (DBI)
1997-11-05 (SMS)
2003-06-27 (SMS)

To the user of this abstract: you may use the entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.