

**ARIZONA GAME AND FISH DEPARTMENT  
HERITAGE DATA MANAGEMENT SYSTEM**

**Plant Abstract**

**Element Code:** PDASC020Z0

**Data Sensitivity:** No

**CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE**

**NAME:** *Asclepias lemmonii* Gray  
**COMMON NAME:** Lemmon's milkweed, bigleaf milkweed  
**SYNONYMS:**  
**FAMILY:** Asclepiadaceae

**AUTHOR, PLACE OF PUBLICATION:** A. Gray, Proceedings of the American Academy of Arts and Sciences 19: 85. 1883.

**TYPE LOCALITY:** Syntype GH-76136: Rucker Valley, Arizona, U.S.A. 1881. Syntype GH-76135: Near Fort Huachuca, Tanner's Canyon, Arizona, U.S.A. Aug 1882.

**TYPE SPECIMEN:** HT: CAS? J.G. Lemmon s.n., 1881. Syntype: GH, US.

**TAXONOMIC UNIQUENESS:** There are 26 species of *Asclepias* in Arizona, and 76 in North America. *Asclepias* is an American and African genus with about 150 species, mostly in the United States and Mexico (Hickman 1993, Lehr 1978, Sundell 1994).

**DESCRIPTION:** Stout perennial deciduous herb. Stems erect, unbranched, 40-150 cm (1.3-5 ft) tall, shaggy pilose. Leaves opposite, subsessile, petioles 2-5 cm long, the blades ovate-elliptic, 6-25 cm (2.4-10 in) long, 3-16 cm (1.2-6.3 in) broad, rounded to shallowly cordate and more or less clasping at the base, rounded to truncate or retuse at the apex, apiculate, pilose to pubescent beneath, sparsely so above, the margins ciliate. Umbels subterminal at the uppermost nodes, 5-8 cm (2-3.2 in) broad, the peduncles 3-12 cm (1.2-4.7 in) long. Flowers large; calyx lobes 4-5 mm long; corolla green to purplish outside, greenish inside, the lobes 9-11 mm long; hoods pale pink, erect-spreading, oblong-quadrate in the lower portion, more or less recurved-divergent at the middle, attenuate to the apex, 7-8 mm long along the dorsal surface, 5-8 mm broad at the top, 4-6 mm longer than the gynostegium (crown of united stamens in milkweeds), the horns radially flat, attached at the base of the hoods, oblong, erect, exserted approximately 1 mm; anther wings 2.5-2.6 mm long; corpusculum (central part of a pollinarium, which is the whole structure that bears the pollinias) 0.4-0.5 mm long, the pollinia 1.6 mm long. Martin et al. (1998) reports flowers cream turning pale yellow. Follicles erect on deflexed pedicels, 9-12 cm long (Sundell 1994).

**AIDS TO IDENTIFICATION:** The following combination of characteristics will help you identify *Asclepias lemmonii*: **1.** Stems are very tall (up to 1-1.5 m), and persistently leafy. **2.** Leaves are very large, opposite, subsessile; blades ovate-elliptic, to 25 cm long and 16 cm

broad, rounded to truncate or retuse at the apex. **3.** Stems are shaggy-pilose, herbaceous above the rootstock. **4.** Hoods are 7-8 mm long, 4-6 mm longer than the gynostegium (Kearney and Peebles 1951, Sundell 1994).

**ILLUSTRATIONS:** Color Photo (Rickett 1970).  
Color photo of Syntype (US-47696, *in* <http://ravenel.si.edu/botany/types/fullRecords.cfm?myFamily=>)  
Color photo of collection (ASU-213423, *in* <http://seinet.asu.edu/collections/TaxaDetails.jsp?>)

**TOTAL RANGE:** Southern and southeastern Arizona south into Chihuahua and Sonora, Mexico.

**RANGE WITHIN ARIZONA:** Cochise County: Huachuca and Chiricahua mountains; Pima County: Baboquivari and Santa Rita mountains; Santa Cruz County: Santa Rita Mountains.

## **SPECIES BIOLOGY AND POPULATION TRENDS**

**GROWTH FORM:** Perennial deciduous forb.

**PHENOLOGY:** Flowering June – September (Aug-Sep per Martin et al., 1998); fruiting August - October.

### **BIOLOGY:**

**HABITAT:** In southeastern Arizona this species is found in the mountains, where it occupies canyons, roadsides, and open woodlands (Sundell 1994). In Mexico, this species is known from open dry slopes, burned areas, and grazed grassland (Martin et al. 1998).

**ELEVATION:** Approximately 5,050 - 7,200 feet (1540-2196 m) in Arizona. Specimens from Mexico have been collected at about 3,934 - 6,885 feet (1200-2100 m), (Martin et al. 1998).

**EXPOSURE:** Often on north and east-facing slopes in Arizona, but has been collected from west-facing slopes. Plants have been found in shaded or partially shaded situations as well as open forest habitats.

**SUBSTRATE:** Definitely known from limestone substrates (Shreve and Wiggins 1964). Probably found on volcanic and other substrates as well, based on known collection locations.

**PLANT COMMUNITY:** Madrean evergreen, pine-oak, and oak woodland communities. One collection (2002, ASU-242979) in mesquite grassland on alluvial outwash from east drainage slope of Huachuca Mts. Associated species include: *Acer* sp. (maple), *Agave*

*palmeri* (Palmer's agave), *A. parryi* var. *huachucensis* (Huachuca agave), *Bouteloua curtipendula* (side-oats grama), *Bouvardia ternifolia* (Scarlet Bouvardia), *Cercocarpus montanus* (Colorado birchleaf mountain-mahogany), *Geranium* sp. (crane's-bill), *Juniperus deppeana* (Alligator juniper), *Oxalis* sp. (woodsorrel), *Pinus cembroides* (Mexican pinyon), *P. discolor* (border pinyon), *P. engelmannii* (Engelmann pine), *P. leiophylla* (Chihuahuan pine), *Platanus racemosa* (California sycamore), *Quercus arizonica* (Arizona oak), *Q. hypoleucoides* (silver-leaf oak), *Q. oblongifolia* (Mexican blue oak), and *Tradescantia* (spiderwort). (Brown 1994, SEINet accessed 2006).

In Mexico, collected in pine/oak woods on sedimentary soil, with *Pinus engelmannii*, *P. chihuahuana*, and *Quercus durifolia* (Cusi barril). (SEINet, accessed 2006).

**POPULATION TRENDS:** Unknown.

## **SPECIES PROTECTION AND CONSERVATION**

**ENDANGERED SPECIES ACT STATUS:** None.  
**STATE STATUS:** None.  
**OTHER STATUS:** Forest Service Sensitive (USDA, FS Region 3 1999)

**MANAGEMENT FACTORS:** Although this species is rather uncommon where it occurs, it does not seem to be very intolerant of disturbance.

**CONSERVATION MEASURES TAKEN:** This species occurs on The Ramsey Canyon Preserve, a protected area. Other known locations in Arizona are all on public or federal lands.

**SUGGESTED PROJECTS:** Areas with known populations should be re-surveyed to determine population trends. The location in the Santa Rita Mountains should be re-verified.

**LAND MANAGEMENT/OWNERSHIP:** DOD – Fort Huachuca Military Reservation; USFS - Coronado National Forest; TNC – Ramsey Canyon Preserve; Private.

## **SOURCES OF FURTHER INFORMATION**

### **REFERENCES:**

- Brown, D.E., ed. 1994. Biotic communities: southwestern United States and northwestern Mexico. University of Utah Press. Salt Lake City, Utah. 342 pp.
- Harvard University Herbaria. 2001. Index of Botanical Specimens. Accessed: 4/26/2006. <http://brimsa.huh.harvard.edu/cms-wb/specimens.jsp>.
- Hickman, J.C., ed. 1993. The Jepson manual, higher plants of California. University of California Press. Berkeley, California. P. 1172.

- Integrated Taxonomic Information System (ITIS). Retrieved 4/26/2006 from ITIS, <http://www.itis.usda.gov>.
- Kearney, T.H., R.H. Peebles with collaborators. 1960. Arizona flora. Second edition with supplement by J.T. Howell, E. McClintock and collaborators. 1951. University of California Press. Berkeley, California. Pp. 655-662.
- Lehr, J.H. 1978. A catalogue of the flora of Arizona. Desert Botanical Garden, Phoenix, Arizona. Pp. 119-120.
- Martin, P.S., et al. 1998. Gentry's Rio Mayo Plants: the tropical deciduous forest and environs of northwest Mexico. The University of Arizona Press. Tucson, Arizona. P. 214.
- Missouri Botanical Garden – TROPICOS, Nomenclatural Data Base. *Asclepias lemmonii* A. Gray. <http://mobot.mobot.org/>. Accessed: 26 Apr 2006.
- NatureServe. 2006. NatureServe Explorer: An online encyclopedia of life [web application]. Version 4.7. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: April 26, 2006).
- Rickett, H.W. 1970. Wildflowers of the United States, Volume four: Part two of three parts: The southwestern states. McGraw-Hill Book Company. New York. Pp. 436-437.
- SEINet. Collections Search Result. Accessed 4/26/2006 at <http://seinet.asu.edu/collections/listDisplay.jsp>.
- Shreve, F. and I.L. Wiggins. 1964. Vegetation and flora of the Sonoran Desert: Volume two. Stanford University Press. Stanford, California. P. 1112.
- Sundell, E. 1994. A new flora for Arizona in preparation: Asclepiadaceae. Journal of the Arizona-Nevada Academy of Science 27(2): 169-187.
- U.S. National Herbarium Type Specimen Register (US). <http://ravenel.si.edu/botany/types/fullRecords.cfm?myFamily=>. Accessed: 9/21/2005.
- USDA, Forest Service Region 3. 1999. Regional Forester's Sensitive Species List.
- USDA, NRCS. 2004. The PLANTS Database, Version 3.5 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

**MAJOR KNOWLEDGEABLE INDIVIDUALS:**

- Frank Reichenbacher - Southwest Field Biologists, Tucson, AZ.  
Eric Sundell - University of Arkansas at Monticello, Monticello, AR.

**ADDITIONAL INFORMATION:**

The genus *Asclepias* was named after Asklepios, a legendary Greek physician and god of medicine.

See Sundell (1994), pg. 170, for a drawing of a typical *Asclepias* flower with the flower parts labeled.

**Revised:** 1999-10-27 (DJG)  
1999-10-28 (DJG)  
2006-05-04 (SMS)

To the user of this abstract: you may use the entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.