

**ARIZONA GAME AND FISH DEPARTMENT
HERITAGE DATA MANAGEMENT SYSTEM**

Animal Abstract

Element Code: ARADE02120

Data Sensitivity: No

CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE

NAME: *Crotalus viridis* Rafinesque (1818)
COMMON NAME: Green Prairie Rattlesnake; Prairie Rattlesnake
SYNONYMS: *Crotalinus viridis viridis* (Rafinesque 1818)
FAMILY: Viperidae

AUTHOR, PLACE OF PUBLICATION: Rafinesque (1818), Amer. Month. Mag. Crit. Rev., Vol. 4, No. 1, Nov. p. 41. Also in Crothers et al. 2008. Squamata: Snakes. In B. I. Crother (ed.), Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico, pp. 46-65 SSAR Herpetological Circular 37.

TYPE LOCALITY: “The type locality was stated as “the Upper Missouri [Valley]” although it was later restricted to “Gross, Boyd County, Nebraska” by Smith and Taylor (1950).” (Degenhardt et al. 1996).

TYPE SPECIMEN: No type specimen was designated.

TAXONOMIC UNIQUENESS: The family Viperidae, commonly known as pit vipers, is a medium to large sized group. Only one of the three subfamilies, Crotalinae, actually penetrates into North America and it contains approximately 140 constituents. Based on Crothers et al. (2008), “Pook et al. (2000, Mol. Phylog. Evol. 15: 269–282), Ashton and de Queiroz (2001, Mol. Phylog. Evol. 21: 176–189), and Douglas et al. (2004, Biology of the Vipers, Schuett, Hoggren, Douglas, Greene [eds.] Eagle Mountain Press) analyzed mtDNA sequence data and concluded that *Crotalus viridis* comprised at least two clades, *C. viridis* and *C. oregonus*, with *C. cerberus* being the sister taxon to populations of *C. oregonus*. The former two studies did not formally recognize *C. cerberus* as a species, although both suggested that it was an evolutionary species based on sequence differences and allopatry. The latter study did recognize *C. cerberus* as well as four other taxa. We take the conservative action supported by the congruence among all three studies, which is the recognition of *C. viridis*, *C. oregonus* and *C. cerberus*.” In addition, *Crotalus viridis* (Rafinesque, 1818) the Prairie Rattlesnake, synonymized *C.v. nuntius* with *C. v. viridis* (Douglas et al. 2004, Biology of the Vipers, Schuett, Hoggren, Douglas, Greene [eds.] Eagle Mountain Press).

DESCRIPTION: The snout-vent length of is commonly greater than 84 cm (33 in) and may reach up to 162 cm (64 in). This snake may present highly variable ground color throughout its range spanning from greenish-gray to yellow, tan or brown (Brennan 2006; Degenhardt et al. 1996; IMNH 1997; Stebbins 2003); more often is tan with brown dorsal blotches. Their

bodies are patterned with 35-55 dark brown blotches on the body (excluding the tail) that are bordered by white, with two rows of smaller spots on their sides. The belly is white and contains no spots (Degenhardt et al. 1996). Typically 27 or 25 dorsal scale rows at midbody; the dorsal scales are keeled. A light stripe extends from behind the eye to the corner of the mouth (Stebbins 2003) and two diagonal white lines can be seen on the side of the head (Degenhardt et al. 1996). The pupils are vertically elliptical. The head is broad and triangular and the neck is slender. It has four or more internasals scales in contact with the rostral; and the anal plate is entire. As with other members of the genus, a rattle composed of a series of loosely interlocking keratinous sections sits on the end of the tail. A new section is added each time the snake sheds its skin. (Brennan 2006).

AIDS TO IDENTIFICATION: *C. viridis* can be distinguished from similar species by the presence of more than two internasals touching the rostral (Ashton & de Queiroz; Brennan & Holycross 2006; Degenhardt et al. 1996; Stebbins, 2003). They may resemble *Sistrurus catenatus* (Massasauga Rattlesnake) which has nine symmetrical plates on the top of its head instead of many more and smaller scales (Degenhardt et al. 1996; TMM@UTA 2000). In addition, *S. catenatus* is smaller (47-76 cm [18-30 in]) than *C. viridis*.

ILLUSTRATIONS:

Color photo (Gary Nafis, CaliforniaHerps.com, Accessed 01/02/2007)

Color illustration (Stebbins 2003: Plate 51)

Color photo (Degenhardt et al. 1996: Plate 121)

Color photo (Bill Sloan, in Arizona Herpetological Association,

<http://www.sloanmonster.com/images/PrarieRtlr.jpg>, Accessed 01/02/2007)

Color photo (Roger A. Repp, in Tucson Herpetological Society

<http://www.arts.arizona.edu/herp/CRVI.html>, Accessed 01/02/2007)

Color photos (Randy Babb, in Brennan 2006 at <http://www.reptilesfaz.com/Snakes-Subpages/h-c-viridis.html>, Accessed 01/02/2007)

Color photo (Suzanne L. Collins 2001, in CNAH <http://www.cnah.org/detail.asp?id=96>)

Color photo (Frank C. Johnson, Accessed 3/28/2007 from

<http://dwrcdc.nr.utah.gov/rsgis2/Search/Display.asp?F1Nm=crotvivi>)

TOTAL RANGE: *C. viridis* has a large distribution from southern Canada to Mexico, east throughout Nebraska, and much of Kansas, and west as far as Idaho in the north, and northeastern Arizona in the south.

RANGE WITHIN ARIZONA: *C. viridis* occurs in eastern and portions of northern Arizona in Apache, Navajo and Coconino counties (Brennan 2006).

SPECIES BIOLOGY AND POPULATION TRENDS

BIOLOGY: Active during the months of March through November, *C. v. viridis* is primarily a

nocturnal ground dweller, but is crepuscular during cooler seasons; they will occasionally climb into trees and shrubs. They are diurnal in the spring and on mild or overcast summer days. In late fall and winter, *C. v. viridis* hibernates alone or in a group den in such structures as rodent burrows (e.g. prairie dog towns), slumped blocks and rock escarpments (Watson & Russell 1997); they may form dens with other snakes including *Coluber constrictor* (Racer), *Pituophis melanoleucus* (Gopher Snake), *Lampropeltis triangulum* (Milk Snake), and *Thamnophis elegans* (Western Terrestrial Garter Snake) (BISON-M 2007). (Degenhardt et al. 1996; Brennan 2006; Brennan & Holycross 2006). In Utah, the species has been observed to move in the summer up to about 1.5 km from the winter den (Hirth et al. 1969, in NatureServe 2006).

Like the other “pit-vipers” (members of subfamily Crotalinae) this snake uses heat sensing pits (one on each side of the face between the eye and nostril) to detect warm-blooded predators and prey (Brennan 2006). Predators include *C. constrictor* (Eastern Racer), *Aquila chrysaetos* (Golden eagle) (BISON-M 2007), various mammalian carnivores, raptors, whipsnakes and Kingsnakes (NatureServe 2006). The rattle is a series of horny rings formed of keratin. These grate against each other in pulses to cause the rattle, the rate depending on temperature and humidity. The pulsing cycle may be 21 per second at 50 °F, 99 per second at 104 °F. The rattle begins with a soundless button on a young snake and grows with age. (BISON-M 2007).

REPRODUCTION: Sexual maturity of male *C. v. viridis* usually occurs at the age of 3-4 years, while females mature at the age of 5-7 years (Watson & Russell 1997): females are live-bearing. In New Mexico, spermatogenesis is known to occur during mid to late summer, from June through August or into September (Degenhardt et al. 1996). In Arizona, *C. viridis* mate in June and July (Brennan 2006) and in southern Alberta, Canada *C. v. viridis* are also known to mate during mid to late summer and into early fall (Watson & Russell 1997). However, it has been found in Nebraska that *C. v. viridis* may undergo two periods of mating—the first in early spring and the second in mid to late summer (Degenhardt et al. 1996). Females will store sperm over the winter and give birth the following year. Spring migrations are known to take place in the northern populations of *C. v. viridis*, where pregnant females may form rookeries a short distance from the hibernacula (Graves & Duvall 1993), while males have been known to migrate long distances in search of foraging areas that contain abundant supplies of food (Duvall, et al. 1985; Duvall, et al. 1989). In Arizona, birthing occurs in the summer months (Brennan 2006), while in Canada females will give birth from late August through mid October. Litter size has been found to be highly correlated with body size and each female may produce 5-14 offspring per clutch (Degenhardt et al. 1996). Young average 24 cm (9.5 in) in length. Females may not produce young each year, although it can occur, rather it is thought that *C. v. viridis* follow a biennial or triennial reproductive cycle (Degenhardt et al. 1996; Watson & Russell 1997).

FOOD HABITS: *C. viridis* will consume birds and their eggs, lizards, and amphibians, but adults appear to prefer small mammals such as mice and other rodents (Brennan 2006;

Degenhardt et al. 1996). Members of the family Viperidae are known as pit vipers because they have a bladder in the back of their head that contains venom, which they use to paralyze prey, but it can also serve as a defense mechanism. The venom is delivered by small grooves in retractable fangs that act as a syringe to inject the neurotoxin into its victim.

HABITAT: As their common name suggests, *C. v. viridis* choose to inhabit remnant or short-grass prairie grasslands, but may also occur in open desert or shrublands, rocky slopes, and montane ponderosa pine forests (Ashton & de Queiroz 2001; Degenhardt et al. 1996; Watson & Russell 1997).

ELEVATION: In Arizona, *C. viridis* can be found from 5,000 feet to at least 9,000 feet (Brennan 2006; Degenhardt et al. 1996). Based on UAZ collection records (UAMNH 2006), *C. v. viridis* ranges from 5,660 to 7,100 feet (1725-2164 m).

PLANT COMMUNITY: Various, but generally with a grassland component as the common name of this snake would imply.

POPULATION TRENDS: Unknown. The subspecies *viridis* is restricted to eastern portion of state in the vicinity of Springerville and Saint Johns. Little work has been done on current distribution, thus trend information is lacking.

SPECIES PROTECTION AND CONSERVATION

ENDANGERED SPECIES ACT STATUS: None
STATE STATUS: None
OTHER STATUS: PR (Mexican Federal Endangered Species List, 2000)

MANAGEMENT FACTORS: Unknown.

PROTECTIVE MEASURES TAKEN:

SUGGESTED PROJECTS: Distribution, trend, and habitat use information is lacking in Arizona.

LAND MANAGEMENT/OWNERSHIP: State Land Department; Private.

SOURCES OF FURTHER INFORMATION

REFERENCES:

Ashton, K.G., A. de Queiroz. 2001. Molecular Systematics of the Western Rattlesnake, *Crotalus*

- viridis* (Viperidae), with Comments on the Utility of the D-Loop in Phylogenetic Studies of Snakes. *Molecular Phylogenetics and Evolution*: Vol. 21, No. 2, pp. 176-189.
- [BISON-M] Biota Information System of New Mexico. 2007/03/28 <http://www.bison-m.org>. Accessed 2007/03/28.
- Brennan, T.C. 2006. Reptiles of Arizona: PRAIRIE RATTLESNAKE, *Crotalus viridis*. Accessed December 29, 2006 at <http://www.reptilesfaz.com/Snakes-Subpages/h-c-viridis.html>.
- Brennan, T.C., A.T. Holycross. 2006. A Field Guide to Amphibians and Reptiles in Arizona. Arizona Game and Fish Department Publication. Phoenix, Arizona. P. 130.
- Crother, B.I. et al. 2003. Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico: Update. J.J. Moriarty, Editor, SSAR, *Herpetological Review* 34(3), 201.
- Crothers et al. 2008. Squamata: Snakes. In B. I. Crother (ed.), *Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico*, pp. 46-65 SSAR *Herpetological Circular* 37.
- Degenhardt, W.G., C.W. Painter, and A.H. Price. 1996. *Amphibians & Reptiles of New Mexico*. University of New Mexico Press. Albuquerque, New Mexico. Pp. 351-353.
- Duvall, D., D. Chiszar, W.K. Hayes, J.K. Leonhardt, and M.J. Goode. 1989. Chemical and Behavioral Ecology of Foraging in Prairie Rattlesnakes (*Crotalus viridis viridis*). *Journal of Chemical Ecology*: Vol. 16, No. 1, pp. 87-101.
- Duvall, D., M.B. King, and K.J. Gutzwiller. 1985. Behavioral Ecology and Ethology of the Prairie Rattlesnake. *National Geographic Research*: Vol. 1, No. 1, pp. 80-111.
- Graves, B.M., and D. Duvall. 1993. Reproduction, Rookery Use, and Thermoregulation in Free-ranging, Pregnant *Crotalus v. viridis*. *Journal of Herpetology*: Vol. 27, No. 1, pp. 33-41.
- Idaho Museum of Natural History (1997) *Crotalus viridis* (Western Rattlesnake) [online], available: <http://imnh.isu.edu/digitalatlas/bio/reptile/serp/crvi/crvifram.htm> [accessed 29 Dec 2006].
- NatureServe. 2006. NatureServe Explorer: An online encyclopedia of life [web application]. Version 6.1. NatureServe Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: March 28, 2007).
- Proyecto de NORMA Oficial Mexicana PROY-NOM-059-ECOL-2000 (Mexican Federal Endangered Species List).
- Stebbins, R.C. 2003. A Field guide to Western Reptiles and Amphibians. Third edition. Houghton Mifflin Company. Boston, Massachusetts. Pp. 414-416.
- Texas Memorial Museum at UT Austin. 2000. Herps of Texas – Snakes: *Crotalus viridis*, Western Rattlesnake (online). Accessed: 28 March 2007. <http://www.zo.utexas.edu/research/txherps/snakes/crotalus.viridis.html>.
- The University of Arizona Museum of Natural History. 2006. Collections: Herps [online]. <http://eebweb.arizona.edu/collections/Herp/HerpPage.htm>. Accessed: 4 April 2007.
- Utah Division of Wildlife Resources, State of Utah Natural Resources. Green Prairie (Prairie) Rattlesnake, *Crotalus viridis viridis*. Accessed: 28 March 2007 at <http://dwrcdc.nr.utah.gov/rsgis2/Search/Display.asp?F1Nm=crotvivi>.
- Valle, L. 2000. Animal Diversity Web, *Crotalus viridis* (On-line). Accessed 12/29/2006 at

http://animaldiversity.ummz.umich.edu/site/accounts/information/Crotalus_viridis.html.

Watson, S.M., and A.P. Russell. 1997. Status of the Prairie Rattlesnake (*Crotalus Viridis viridis*) in Alberta. Alberta Environmental Protection, Wildlife Management Division, Wildlife Status Report No. 6, Edmonton, AB. 26pp.

MAJOR KNOWLEDGEABLE INDIVIDUALS:

ADDITIONAL INFORMATION:

Origin of Scientific Name (Beltz 2006):

Crotalus: Greek – krotalon, a rattle – ref. rattle on tail, typical of all but one member of the genus.

viridis: Latin – green – some adults on Great Plains have light greenish-gray dorsal coloration.

** As Brennan (2006) reports, “This rattlesnake is capable of delivering large amounts of potent venom. If encountered it should be left alone. Most envenomations occur when the victim attempts to capture or kill the animal.”

Revised: 2007-01-08 (TFH)
2007-04-06 (SMS)
2010-11-09 (SMS)

To the user of this abstract: you may use the entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.