

**ARIZONA GAME AND FISH DEPARTMENT
HERITAGE DATA MANAGEMENT SYSTEM**

Animal Abstract

Element Code: ABPBR01030

Data Sensitivity: No

CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE

NAME: *Lanius ludovicianus* L.

COMMON NAME: Loggerhead Shrike

SYNONYMS:

FAMILY: Laniidae

AUTHOR, PLACE OF PUBLICATION: Linnaeus, 1766

TYPE LOCALITY:

TYPE SPECIMEN:

TAXONOMIC UNIQUENESS: 21+ Species in genus.

DESCRIPTION: Slightly smaller than the American Robin; length 23.0 cm (9.0 in), weight 48 g (1.7 oz). Head and back bluish-gray; underparts white, very faintly barred. The head is large in proportion to its body size. Broad black mask extends above eye, thinly across top of bill. Birds in California darker than in other parts of range. All dark bill stout, conical with hook. Rump gray to white. Juvenile paler, barred overall with brownish-gray upperparts; acquiring adult plumage by first fall.

AIDS TO IDENTIFICATION: Loggerhead Shrikes are distinguished from Northern Shrikes by their slightly smaller size, 20 cm relative to 25cm. Also, the black mask on the Northern shrike does not extend across the forehead above the beak as in the Loggerhead. The base of the lower mandible is black instead of pale, and the beak on the Northern is about one-half the length of the head (front to back) whereas it is substantially shorter on the Loggerhead. In flight, the tail and wings are darker, and the white wing patches are smaller than in Northern Mockingbird. (National Geographic Society, 1999).

ILLUSTRATIONS: Color Photos (www.birdphotography.com)
Distribution Map (i-bird.com)
Color Photo and Distribution map (Yosef, 1996)
Color drawings (Sibley, 2000: p. 341)
Color drawing (National Geographic, 1999: p. 305)
Color drawing (Peterson, 1990: p. 280)
Color drawing of egg (Baicich 1997)

TOTAL RANGE: Breeds throughout southern Canada, south through Great Basin; in California to Baja California, Mexico, along Gulf coast to southern Florida. Withdraws southward from northern half of breeding range for winter.

RANGE WITHIN ARIZONA: "More or less common summer resident throughout open parts of state (except in brushless grassland) below Transition Zone, rather uncommon (at least in midsummer) along Mexican border west of Baboquivari Mountains. Fairly common transient in Transition Zone at least in fall. Winters commonly in Lower Sonoran Zone, less commonly in open Upper Sonoran Zone even into evergreen oaks" (Monson and Phillips 1981).

SPECIES BIOLOGY AND POPULATION TRENDS

BIOLOGY: Territorial throughout the year. Size of territory may be about 6 ha in grassy hills; 10-16 ha in semidesert. Song is medley of low warbles and harsh squeaky notes. "Song generally sharp, precise, mechanical two-syllable phrases *krrDI* or *JEEuk* (etc.) repeated over and over at short intervals. Calls include a harsh, scolding *jaaa*, grating *teen raad raad raad raad*, and other variations similar to song phrases." (Sibley 2000). Often perches on poles, wires or fence posts. They hunt in open or brushy areas, diving from a low perch, then rising swiftly to the next lookout. They often sit immobile for long periods watching for prey. Lacking the talons of raptors, it stuns or kills flying birds with a blow from its powerful beak. Has habit of impaling prey on thorns either to eat them immediately or to larder them for future use. Shrikes show an amazing memory for the placement of their victims: in Texas, shrikes were reported returning to mummified frogs they had stored eight months before. Species is considered very aggressive in defending territory and both genders participate. Sexes hold separate territories during nonbreeding seasons.

REPRODUCTION: During courtship the male feeds the female and performs a flight display. Mock pursuits also occur. Nests in open areas in forests or in open country. May nest in hedgerows or trees. Nest is bulky, cup-shaped, of twigs lined with rootlets, grasses, string, feathers, deer hairs, 8-15 ft above ground. Both males and females select site and gather materials, but female constructs the nest. Female incubates 4-5, sometimes 6-7 dull white to light gray or buff, spotted eggs for 14-16 days. Incubation by both sexes also reported. The nestlings are altricial and downy with bright orange skin. The down is white and very sparse being on the body only. The bill is buffish-yellow with the mouth and the gape flanges being yellow as well. Two broods per season, young tended by both adults, independent in 36 days.

FOOD HABITS: Only know predatory songbird. Feeds primarily on insects, small birds, lizards and rodents. Diet varies with season and location. Mice may be half of diet in winter. Small birds taken are warblers, sparrows, mockingbirds, bluebirds, swallows and other passerines. Hunt in open or brush areas in a sit and wait strategy, diving from low perch, and then rising swiftly to next lookout. Sometimes referred to as the butcher bird after its habits

of impaling prey on thorns. This facilitates dismembering the prey in as much as shrikes have no talons.

HABITAT: Open country with scattered trees and shrubs, savanna, desert scrub and occasionally open woodland. Often found on poles, wires or fence posts.

ELEVATION: Unknown

PLANT COMMUNITY: Unknown

POPULATION TRENDS: In the past few years, the loggerhead shrike has aroused serious concern because of its declining numbers throughout its range in Canada and the United States. It has been declining in North America since the 1960s. Decline has been recorded in all regions of the country, even those areas with great amounts of habitat. Is considered moderately threatened throughout its range.

SPECIES PROTECTION AND CONSERVATION

ENDANGERED SPECIES ACT STATUS: None (USDI, FWS 1996)
[C2 USDI, FWS 1991]

STATE STATUS: None

OTHER STATUS: None (USDI, BLM AZ 2005)
[Bureau of Land Management Sensitive
(USDI, BLM AZ 2000)]

MANAGEMENT FACTORS: Being high on the food chain, they are subject to pesticide poisoning. Fire exclusion, habitat loss, and dependency on edge habitat which increases predation pressure are also factors that need to be considered for this species.

PROTECTIVE MEASURES TAKEN: Unknown

SUGGESTED PROJECTS: Unknown

LAND MANAGEMENT/OWNERSHIP: Unknown

SOURCES OF FURTHER INFORMATION

REFERENCES:

Baicich, P. J. & Harrison, C. J. 1997. A Guide to the Nests, Eggs, and Nestlings of North American Birds. Natural World Academic Press, San Diego, California. Pp: 265, Pl. 58.
Canadian Wildlife Service, National Site, (http://www.cws-scf.ec.gc.ca/index_e.cfm).

- Ehrlich, P. R. et al. 1988. The Birder's Handbook. Simon & Schuster Inc. New York, New York. Pp: 466-476.
- Monson G. and A.R. Phillips. 1981. Annotated checklist of the birds of Arizona, second edition. The University of Arizona Press, Tucson. p. 118.
- National Geographic. 1999. Field guide to the Birds of North America. Third Addition. National Geographic Society, Washington, D.C. pp. 304-305.
- NatureServe Explorer: An online encyclopedia of life [web application]. 2001 Version 1.6. Arlington Virginia, USA: NatureServe. Available: <http://natureserve.org/explorer>. (Accessed: November 13, 2002).
- Peterson, R.T. 1990. A field guide to western birds, third edition. Houghton Mifflin Company, Boston. pp. 280-281.
- Pruitt Lori, 2000. Loggerhead Shrike Status Assessment, USFWS report, USFWS Bloomington, IN.
- Scott, S.L. 1987. National Geographic Society field guide to the birds of North America, second edition. National Geographic Society, Washington, D.C. p. 334.
- Sibley, D.A. 2000. National Audubon Society, The Sibley Guide to Birds. A Chanticleer Press Edition. Alfred A. Knopf, New York. P. 341.
- Terres, J.K. 1980. The Audubon Society encyclopedia of North American birds. Alfred A. Knopf, New York pp. 819-821.
- USDI, Bureau of Land Management. 2000. Arizona BLM Sensitive Species List. Instruction Memorandum No. AZ-2000-018.
- USDI, Bureau of Land Management. 2005. Arizona BLM Sensitive Species List.
- USDI, Fish and Wildlife Service. 1991. Endangered and Threatened Wildlife and Plants; Animal Candidate Review for Listing as Endangered or Threatened Species, Proposed Rule. Federal Register 56(225):58811.
- USDI, Fish and Wildlife Service. 1996. Endangered and Threatened Wildlife and Plants: Review of Plant and Animal Taxa that are Candidates for Listing as Endangered or Threatened Species. Federal Register 61(40):7596-7613.
- Yosef R. 1996. Loggerhead Shrike (*Lanius ludovicianus*). In The Birds of North America, No. 231 (A. Poole and F. Gill eds.). The Academy of Natural Sciences, Philadelphia, and The American Ornithologists' Union, Washington, D.C.

MAJOR KNOWLEDGEABLE INDIVIDUALS:**ADDITIONAL INFORMATION:**

Revised: 1995-08-08 (DBI)
 1997-03-05 (SMS)
 2002-11-14 (RHB)
 2004-01-02 (AMS)

To the user of this abstract: you may use the entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.