

**ARIZONA GAME AND FISH DEPARTMENT
HERITAGE DATA MANAGEMENT SYSTEM**

Animal Abstract

Element Code: AMACC01070

Data Sensitivity: Yes

CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE

NAME: *Myotis evotis*
COMMON NAME: Long-eared Myotis
SYNONYMS: None
FAMILY: Vespertilionidae

AUTHOR, PLACE OF PUBLICATION: H. Allen 1864.

TYPE LOCALITY:

TYPE SPECIMEN:

TAXONOMIC UNIQUENESS: *Myotis evotis* is one of 88 species of *Myotis* worldwide and 9 *Myotis* species within Arizona. This species is composed of from two to six subspecies based on analysis of cranial morphology and pelage characteristics. The two commonly recognized subspecies are *M. e. evotis* and *M. e. pacificus*. Manning (1993) described 6 subspecies of *M. evotis*, including *M. evotis milleri*, currently recognized as a unique species, *M. milleri*.

DESCRIPTION: Large bat (length of head and body 75.0-100.0 mm (3.0-3.72 in.) with long blackish ears (18.0-23 mm [0.32-0.9 in.] long) extending 5.0-10.0 mm (0.2-0.4 in.) beyond tip of nose when laid flat forward, length of forearm 35.5-41.0 mm (1.42-1.64 in.). Wingspan is 25-30 cm (10-12 in). Adult weight 5-8 g. Fur on upper parts light to medium brown pale brownish to straw-colored overall, full, soft, glossy. About 10.0 mm (0.4 in.) long middorsally with individual hairs black at base. Posterior border of uropatagium with inconspicuous fringe of minute hairs; tragus long and slender. Calcar extending about one half way from foot to tip of tail, not keeled or only slightly so. Cranium rises gradually from rostrum to braincase, skull fairly narrow.

AIDS TO IDENTIFICATION: No conspicuous fringe of hairs on posterior border of the uropatagium. Long tooth rows with robust molars, and auditory bullae relatively large when compared to other long-eared species. Easily confused with *M. auriculus*, which is similar in size, ear length, and color, but tends to have brownish ears and membranes with a more dull, brownish pelage overall.

ILLUSTRATIONS: B&W photo (Burt 1976:26, plate II)
B&W photos (Manning and Jones Jr. 1989:328,329; Fig.1, 2)

Color photo (Peterson *in*
www.enatur.../showSpeciesGS.asp?curGroupID=5&curPageNum=39&re
cnum=MA015)
Color photo (Tuttle *in* http://www.batsnorthwest.org/myotis_evotis.html)
Color photo (Wilson 1999)
Color photo (Harvey 1999)

TOTAL RANGE: Occurs in temperate western North America from central British Columbia and southern Saskatchewan and Alberta southward along the Pacific Coast to Baja California, eastwardly through Montana and Idaho to the western Dakotas, and from Nevada, Utah, Wyoming, and Colorado to New Mexico and Arizona.

RANGE WITHIN ARIZONA: Kaibab Plateau, Mogollon Plateau.

SPECIES BIOLOGY AND POPULATION TRENDS

BIOLOGY: Long-eared Myotis are categorized as "hovering gleaners" which feed by taking prey from the surface of foliage, tree trunks, rocks, or ground. They are fast and maneuverable in flight, have been observed hovering and are often observed foraging in dense vegetation. *M. evotis* varies the echolocation frequencies and patterns used in response to different foraging situations. They typically use lower frequency calls and passive listening in order to detect prey, making them particularly adept at capturing tympanate moths which are sensitive to the typical echolocation frequencies used by insectivorous bats. *M. evotis* seem particularly efficient at foraging in high elevation habitats and when ambient temperatures are low. Unlike most other insectivorous bats, where females are typically restricted to lower elevations and warmer conditions during the breeding season, reproductively active *M. evotis* females do occur throughout the elevational range of this species throughout the breeding season. These bats probably migrate short distances between summer haunts and winter retreats, although very little is known about these migration patterns and nothing is known about hibernacula. It has been suggested that they roost in the winter primarily in caves and abandoned mines. The only known hibernation site is an abandoned mine in Montana. The record longevity for this species is 22 years, but most individuals probably live for much less than this. A snake, the yellow-bellied racer, is a known predator of these bats in British Columbia.

REPRODUCTION: Mating occurs in the fall; probably around the time these bats enter their winter hibernacula. Ovulation and fertilization occur in the following spring. Earliest reported date of pregnancy for this species is May 19 in California, the latest reported pregnancy was July 7 in British Columbia. Females give birth to one young per year in late June or July. Females form small maternity colonies in the summer, while males and barren females live singly or in small groups, occasionally occupying the same roost as the maternity colony but roosting apart from it.

FOOD HABITS: Jameson and Peeters (1988) state that *M. evotis* is "a late flying species, emerging after dark. Forages low from 4-6 ft. above ground". However, other evidence (Hoffmeister, 1986) reports two specimens taken in early evening. Foraging times probably vary with prey availability, ambient temperature, and reproductive status. These bats are often captured when air temperatures are low, but may be more active when insect activity is highest. *M. evotis* feeds on Lepidoptera, Coleoptera, Diptera, Neuroptera, Hymenoptera, Hemiptera, and Homoptera. The majority of prey taken by *M. evotis*, though, is Lepidopterans. Differences in food taken have been observed with males eating primarily moths and females taking more beetles. Where *M. evotis* occurs sympatrically with *M. auriculus*, there is evidence that *M. evotis* of both sexes prey mainly on beetles, while *M. auriculus* individuals prey mainly on moths.

HABITAT: According to Hoffmeister, 1986, *M. evotis* inhabits ponderosa pine or spruce-fir forests of Arizona. During the summer months these bats roost in small groups of 12 to 30 individuals in rock outcroppings, tree cavities, under peeling bark, in stumps, caves, mines, sink holes, lava tubes, or in abandoned buildings. Large diameter trees and snags seem to be the preferred tree roost sites (Rabe, 1998, Waldien et. al., 2000). During winter it is likely that they use caves and abandoned mines as hibernacula. *M. evotis* are most often captured in mixed coniferous forests but also occur in higher elevation forests, pinyon-juniper woodlands, sagebrush steppe, and in riparian desert scrub habitats. The availability of appropriate roost sites may more strongly influence local distribution and abundance than plant community composition. In Oregon, areas where *M. evotis* forage and roost seem to be strongly influenced by the availability of water sources as well. Foraging areas and day roosts were more likely to be found close to a water source and were less influenced by forest composition (Waldien and Hayes, 2001).

ELEVATION: Sea level to 10,000 ft. (3,058 m).

PLANT COMMUNITY: Juniper, spruce-fir, and a few pine.

POPULATION TRENDS: Stable, though unique populations inhabiting relatively isolated mountain ranges may be threatened by loss of habitats. *M. evotis* is moderately common in areas of suitable habitat but may be threatened by loss of suitable roost sites throughout its range.

SPECIES PROTECTION AND CONSERVATION

ENDANGERED SPECIES ACT STATUS: None (USDI, FWS 1996)
[C2 USDI, FWS 1994]

STATE STATUS: None

OTHER STATUS: Bureau of Land Management Sensitive
(USDI, BLM AZ 2000, 2005)

Determined Subject to Special Protection for *M. evotis evotis* (Secretaría de Medio Ambiente 2000)
[Listed Rare for *M. evotis evotis*, Secretaría de Desarrollo Social 1994]

MANAGEMENT FACTORS: The lack of understanding of intra-specific variation within this species compromises the effectiveness of current management policy.

PROTECTIVE MEASURES TAKEN:

SUGGESTED PROJECTS: "Variation in *Myotis evotis* has not been studied since Miller and Allen's (1928) revision of the genus. A thorough morphometric analysis of the species throughout its known range is overdue" (Manning and Jones).

LAND MANAGEMENT/OWNERSHIP: USFS – Coconino, Apache-Sitgreaves and Kaibab National Forests.

SOURCES OF FURTHER INFORMATION

REFERENCES:

- Barclay, R.M.R. 1991. Population structure of temperate zone insectivorous bats in relation to foraging behavior and energy demand. *Journal of Animal Ecology*, 60(1): 165-178.
- Best, T.L. et al. Available:<http://talpa.unm.edu/batcall/accounts/accountsbase/myoevo.html>.
- Bogan, M.A. 1999. *Myotis evotis*. In *The Smithsonian Book of North American Mammals*, D.E. Wilson and S. Ruff, Eds. Smithsonian Institution Press, Washington and London.
- Burt, W.H. 1976. *A field guide to the mammals*. Third edition. Houghton Mifflin Co., Boston. pp.27-29.
- Faure, P.A. and R.M.R. Barclay. 1992. The sensory basis of prey detection by the long-eared bat, *Myotis evotis*, and the consequences for prey selection. *Animal Behavior*, 44(1): 31-39.
- Faure, P.A. and R.M.R. Barclay. 1994. Substrate-gleaning versus aerial hawking: plasticity in the foraging and echolocation behavior of the long-eared bat, *Myotis evotis*. *Journal of Comparative Physiology*, 174(5): 651-660.
- Hall, E.R. and K.R. Kelson. 1959. *The mammals of North America*. Volume I. The Ronald Press Company. pp.168-170.
- Harvey, M. J. et al. 1999. *Bats of the United States*. Arkansas Game and Fish Commission. P. 40.
- Hoffmeister, D.F. 1986. *Mammals of Arizona*. The University of Arizona Press. Tucson, Arizona. p.71.
- [Http://sevilleta.unm.edu/data/species/mammal/socorro/profile/long-eared-myotis.html](http://sevilleta.unm.edu/data/species/mammal/socorro/profile/long-eared-myotis.html).
- Ingles, L.G. 1954. *Mammals of California and its coastal waters*. Stanford University Press. Stanford, California. p.71.

- Jameson, E.W. Jr. and H.J. Peeters. 1988. California mammals. California natural history guides. University of California Press. Berkeley, California. pp.128-129.
- Mammal Species of the World. 1993. Second edition. Edited by Wilson, D.E. and D.M. Reeder. Smithsonian Institution Press. Washington, D.C. pp.207-216.
- Manning, R.W. and J.K. Jones Jr. 1989. *Myotis evotis*. Mammalian Species No. 329. The American Society of Mammalogists. pp.1-5.
- NatureServe Explorer: An online encyclopedia of life [web application]. 2001. Version 1.6. Arlington, Virginia, USA: NatureServe. Available: <http://www.natureserve.org/explorer>. (Accessed: March 6, 2003).
- Peterson, B.M. ENature. Available: www.enatur.../showSpeciesGS.asp?curGroupID=5&curPageNum=39&recnum=MA015
- Rabe, M.J., T.E. Morrell, H. Green, J.C. DeVos, C.R. Miller, Jr. 1998. Characteristics of ponderosa pine snag roosts used by reproductive bats in northern Arizona. *Journal of Wildlife Management* 62(2): 612-621.
- Secretaría de Desarrollo Social. 1994. *Diario Oficial de la Federacion*. p. 27.
- Secretaría de Medio Ambiente. 2000. *Diario Oficial de la Federacion*. p. 44.
- Tuttle, M. Available:http://www.batsnorthwest.org/myotis_evotis.html.
- USDI, Bureau of Land Management. 2000. Arizona BLM Sensitive Species List. Instruction Memorandum No. AZ-2000-018.
- USDI, Bureau of Land Management. 2005. Arizona BLM Sensitive Species List.
- USDI, Fish and Wildlife Service, 1994. Endangered and Threatened Wildlife and Plants; Animal Candidate Review for Listing as Endangered or Threatened Species; Proposed Rule. *Federal Register* 59(219):58986.
- USDI, Fish and Wildlife Service. 1996. Endangered and Threatened Wildlife and Plants: Review of Plant and Animal Taxa that are Candidates for Listing as Endangered or Threatened Species. *Federal Register* 61(40):7596-7613.
- USGS. Available: <http://www.npwrc.usgs.gov/resource/distr/others/nddanger/species/myotevot.htm>.
- Waldien, D.L., J.P. Hayes, and E.B. Arnett. 2000. Day roosts of female long-eared *Myotis* in western Oregon. *Journal of Wildlife Management*, 64(3): 785-796.
- Waldien, D.L. and J.P. Hayes. 2001. Activity areas of female long-eared *Myotis* in coniferous forests in western Oregon. *Northwest Science*, (75) 3: 307-31.
- Wilson D.E. et al. 1999. *The Smithsonian Book of North American Mammals*. Smithsonian Institution Press, Washington, in association with the American Society of Mammalogists, pp 88-90.

MAJOR KNOWLEDGEABLE INDIVIDUALS:

- J.K. Jones Jr. - Texas Tech University, Lubbock, Texas.
- R.W. Manning - Texas Tech University, Lubbock, Texas.
- Dr. Michael A. Bogan – USGS – Biological Resources Division and University of New Mexico, Department of Biology.

ADDITIONAL INFORMATION:

Myotis is derived from two Greek words, mys, “mouse” and otos, “ear” in reference to the fact that the ears of many common bats resemble those of mice. *Evotis* means “good ears” and pertains to the conspicuous ears of this bat and is of Greek origin.

Revised: 1992-01-03 (JSP)
1994-03-29 (DBI)
1994-04-07 (DCN)
1997-03-04 (SMS)
2002-05-31 (TAD)
2003-03-08 (AMS)

To the user of this abstract: you may use the entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.