

**ARIZONA GAME AND FISH DEPARTMENT
HERITAGE DATA MANAGEMENT SYSTEM**

Invertebrate Abstract

Element Code: IMGASC9370

Data Sensitivity: No

CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE

NAME: *Sonorella magdalenensis* (Stearns, 1890)

COMMON NAME: Sonoran talussnail

SYNONYMS: *Helix magdalenensis* Stearns, 1890; *Sonorella tumamocensis* Pilsbry & Ferriss 1915; *S. sitiens arida* Pilsbry & Ferriss 1915; *S. hinkleyi* Pilsbry & Ferriss 1919; *S. hinkleyi fraterna* Pilsbry & Ferriss 1919; *S. tumacacori* Pilsbry & Ferriss 1919; *S. cayetanensis* Pilsbry & Ferriss 1919; *S. linearis* Pilsbry & Ferriss 1923

FAMILY: Helminthoglyptidae

AUTHOR, PLACE OF PUBLICATION: (R.E.C. Stearns, 1890, as *Helix*, U.S. Nat. Mus 13(813):205-225; Pls. 15-17).

TYPE LOCALITY: Sonora, Mexico, on top of a mountain, 1,000 ft above Magdalena; elevation not given. Found in 1965 at the probable T.L. in Sierra Magdalena, ca 1 mi N of Magdalena, at 3,650 ft (which is ca 1,000 ft above the town). (Bequaert & Miller, 1973).

TYPE SPECIMEN:

TAXONOMIC UNIQUENESS: Based on an unpublished revision by W.B. Miller (1968a, in Bequaert and Miller 1973), he recognized 68 valid species of *Sonorella* (with 19 subspecies), 57 of them in Arizona (three common with Sonora), 3 in New Mexico, 1 in trans-Pecos Texas (in common with New Mexico), 8 in Sonora (3 in common with Arizona), and 3 in Chihuahua. *Sonorella magdalenensis* is 1 of 23 species in the *S. granulatissima* Complex.

DESCRIPTION: Snails in the genus *Sonorella* have a “depressed globose, helicoids shell, 12 to 30 mm in diameter, umbilicate or perforate, with a wide, unobstructed mouth and a thin, barely expanded peristome, smoothish or slightly sculptured with growth-lines, occasionally with fine oblique or spiral granulation and short hairs (mainly on the early whorls), lightly colored, and normally with a dark peripheral band. Its most characteristic features are, however, in the genitalia, which lack a dart sac and mucus glands.” (Bequaert and Miller, 1973). For species in the *S. granulatissima* Complex: The verge of the penis is usually stout and truncate, reaching extremes of diminution in some species or gigantism in others. Snails in the complex have minutely granulose or wrinkly-granulose shells, with a readily peeling periostracum; mostly without apical spirally descending threads. (Bequaert and Miller, 1973).

AIDS TO IDENTIFICATION: The most characteristic features of the genus *Sonorella* are, in the genitalia, which lack a dart sac and mucus glands (Bequaert and Miller 1972).

ILLUSTRATIONS:

TOTAL RANGE: In Arizona from Pima County and Santa Cruz counties. Sonora, Mexico, from Sierra Magdalena (Type Locality) to as far south as Sierra Pajaritos.

RANGE WITHIN ARIZONA: In Pima County, known from Cerro Colorado; Roskruge Mts.; S part of Tucson Mts.; N foothills of Santa Rita Mts.; and Tumamoc Hill near Tucson. In Santa Cruz County, known from San Cayetano and Tumacacori mountains.

SPECIES BIOLOGY AND POPULATION TRENDS

BIOLOGY: Terrestrial gastropods do not move much, usually only to find food or reproduce. Olfaction is the primary sensory behavior utilized to find and move toward a food item (on the scale of centimeters to meters). A moving terrestrial gastropod lays down water-laden mucus on which it moves, exposing its integument to a potentially drying atmosphere, and increasing its water losses through the pallial cavity because of the necessity for gas exchange. A roosting terrestrial gastropod deploys a variety of passive mechanisms for water conservation, including the direct protection of its wet surfaces from drying conditions, avoidance of temperature extremes, the creation of more favorable microclimates and decreases in gas exchange. (A. Cook, *in* Barker 2001).

REPRODUCTION:

FOOD HABITS: Probably omnivorous, feeding on plant material (including algae, mosses, lichens, and possibly roots, shoots, leaves, flowers, anthers, pollen, fruit, seeds and rotting wood), and microorganisms associated with live and decaying vegetation; followed to a lesser extent by fungi and soil. (Speiser, *in* Barker, 2001).

HABITAT: The talussnail is a rock snail usually found in taluses or "slides" of coarse broken rock, generally found in crevices one to several feet below the surface, sealed to stones by their mucus (SDCP).

ELEVATION: In Arizona, from as low as 2,750 feet up to 6,000 feet (839-1830 m) elevation (Bequaert & Miller, 1973).

PLANT COMMUNITY:

POPULATION TRENDS: Unknown.

SPECIES PROTECTION AND CONSERVATION

ENDANGERED SPECIES ACT STATUS: None
STATE STATUS: 1C (AGFD SWAP 2012)
OTHER STATUS: Forest Service Sensitive (USDA, FS Region 3 2013)

MANAGEMENT FACTORS: Threats include destruction or disturbance of talus slopes.

PROTECTIVE MEASURES TAKEN:

SUGGESTED PROJECTS: Validity of the informal *Sonorella* “species-groups” (or “complexes”) has been brought into question by Naranjo-García (1988) and Roth (1996). Further research, including the use of molecular techniques, is needed to help clarify the relationships of these informal taxa. (Gilbertson and Radke 2005).

LAND MANAGEMENT/OWNERSHIP: USFS – Coronado National Forest; State Land Department; Private.

SOURCES OF FURTHER INFORMATION**REFERENCES:**

- Arizona Game and Fish Department. 2012. Arizona’s State Wildlife Action Plan 2012-2022. Phoenix, AZ.
- Bequaert, J.C., and W.B. Miller. 1973. The Mollusks of the Arid Southwest. The University of Arizona Press. Tucson, Arizona. Pp. 111, 121-122.
- [Http://www.co.pima.az.us/cmo/sdcp/sdcp2/fsheets/vuln/ts.html](http://www.co.pima.az.us/cmo/sdcp/sdcp2/fsheets/vuln/ts.html). Fact-sheet: Talussnail, *Sonorella*. Accessed: 11/23/2005.
- Integrated Taxonomic Information System (ITIS). Retrieved 3/26/2008 from ITIS, <http://www.itis.usda.gov>.
- NatureServe. 2008. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.0. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: March 26, 2008).
- Santa Barbara Museum of Natural History. SBMNH: Invertebrate Zoology Online Database Collection. <http://www.sbcollections.org/iz/recordview.php>. Accessed: 3/26/2008.
- The Field Museum. 2007. Department of Zoology Invertebrates Collections. Accessed 3/26/2008. <http://emuweb.fieldmuseum.org/iz/MolluscDisplay.php>.
- USDA, Forest Service Region 3. 2013. Regional Forester's List of Sensitive Animals.

MAJOR KNOWLEDGEABLE INDIVIDUALS:**ADDITIONAL INFORMATION:**

The genus *Sonorella* occurs over most of Arizona (except a strip north of the Grand Canyon, an extensive northeast corner, and the small southwest *Eremarionta* area), the southwest corner of New Mexico, trans-Pecos Texas, northeast Sonora, and the northwest corner of Chihuahua, Mexico. (Bequaert and Miller, 1973).

Revised: 2008-04-03 (SMS)

To the user of this abstract: you may use the entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.